Документ подписан простой электронной подписью Информация о владельце: ФИО: Косенок Сергей Михайлович Должность: ректор Дата подписания: 10.06.2024 08:16:27 Уникальный программный ключ: **Test task for diagnostic testing in the discipline:** e3a68f3eaa1e62674b54f4998099d3d6bfdcf836

Discipline name, semester: Outpatient therapy, 10th semester, 5th year

Curriculum	31.05.01
	General Medicine
Qualification	General Medicine
Form of education	Full-time
Department	Internal diseases
Graduating department	Internal diseases

Test task for diagnostic testing in the discipline:

Competence being tested	Exercise	Answer options	Question difficulty type
GC-1 GPC- 1 PC-1 PC - 2 PC -7 PC - 11	<i>Indicate one correct answer</i> 1. THE OPTIMAL METHOD OF IDENTIFYING PERSONS AT RISK FOR THE DEVELOPMENT OF A DISEASE IS	a) recording of appealsb) recording of cases oftemporary disabilityc) medical examinationd) all of the above	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate one correct answer.2. For the period of inpatient examination, a contract sentried by the medical commission of the military committee is issued	 a) Certificate of incapacity for work issued by the clinic b) Certificate of incapacity for work issued by the hospital c) Certificate f - 095/u d) Certificate of any form issued by the hospital e) Certificate of any form issued by the clinic 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate one correct answer.3. The WHO criteria for chronic bronchitis is the duration of cough	 a) at least 6 months a year for 2 consecutive years b) more than 4 months in a given year c) at least 3 months a year for 2 years in a row d) at least 2 months a year for 3 years in a row 	low

GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate one correct answer.</i>4. The triad of symptoms most characteristic for chronic bronchitis include	 a) sputum production b) cough c) pulmonary hypertension d) chest pain d) shortness of breath e) sputum production, cough, shortness of breath 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate one correct answer.</i> 5. The main clinical signs of bronchial obstruction are	 a) poor coughing up sputum b) expiratory nature of shortness of breath c) signs of pulmonary emphysema d) bilateral crepitant rales e) "singing" dry wheezing e) coughing up sputum, shortness of breath, signs of emphysema, dry wheezing 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate one correct answer. 6 When you are referred to a sanatorium, a certificat of disability is issued for the whole term of treatment and for the days of travel to the place of treatment and back	 a) Persons affected by the accident at the Chernobyl nuclear power plant b) Liquidators of the consequences of the accident at the Mayak enterprise c) To the internationalist warriors d) All the above categories of citizens 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Укажите правильные ответы 7. АТРОВЕНТ ПРИ ХОБЛ НАЗНАЧАЮТ ПОИНГАЛЯЦИИРАЗА В ДЕНЬ. 1. 1 2. 2 3. 3 4. 4	 А. Правильные ответы 2 и 4 Б. Правильные ответы 3 и 2 В. Правильные ответы 2 и 2 Г. Правильные ответы 1 и 4 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate all correct answers</i> 8. The list of accounting documents for physical checkpoints is not included	a) F. 025/y 6) F. 030/y B) F. 131/y r) F. 095/y	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Please indicate one correct answer. 9. In the event of fever of unexpected genesis, it is necessary to exclude the following infectious diseases: 1. typhoparasitic; 2. typhus, yersiniosis; 3. malaria, hiv; 4. sepsis, tuberculosis. 	 if answers 1, 2 and 3 are correct; if answers 1 and 3 are correct; if answers 2 and 4 are correct; if the correct answer is 4; if answers 1, 2, 3 and 4 are correct. 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate the correct answer 10. The most informative objective clinical sign for diagnosis of lgk lysed form of bronchiectatic disease is	a) shortening of percussiontoneb) hard breathingc) dry wheezing	average

		d) fine bubbling moist ralese) lokalized moist medium- and large-bubble rales	
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate all correct answers 11. An early clinical symptom of central lung cancer is:	 dry cough or cough with mucous sputum, sometimes streaked with blood pulmonary hemorrhage shortness of breath and weight loss fever hemoptysis 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Please indicate all correct answers 12. Detection of erosions in the antral of the stomach during fibrogastroduodenscopy is characteristic for which treatment is carried out 1. granulomatous gastritis 2. diseases of menetrier 3. autoimmune gastritis 4. helicobacter infection 5. gks 6. proton pump inhibitors 7. gastroprotectors 8. eradication therapy 9. udhc 	A. Correct answers 1, 6 and 7 B Correct answers 2, 7 and 9 B. Correct answers 3, 5 and 6 D. Correct answers 4 and 8	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate all correct answers</i> 13. Acute laryngitis in dynamics threatens the development of:	 persistent dysphonia; acute stenosis of the larynx; asthmatic bronchitis; hypersensitivity of the larynx. 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Please indicate all correct answers 14. For the diagnosis of chronic gastritis, pressures: 1. collection of complaints and anamnesis is enough 2. ultrasound of the obp; 3. morphological study of the gastric mucosa (gut); 4. identification of adr; 5.esophagogastroduodenoscopy (egds); 6. ph-metry of gastric contents. 	A. Correct answers 1 and 4 B. Correct answers 3, 4 and 5 B. Correct answers are 2, 4 and 6 D. Correct answers 1, 2 and 4	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate all correct answers</i> 15. Patients suffering with bronchiectatic disease usually present the following complaints: 1. fever, chest pain; 2. cough with easy mouth, often in the morning; 3. hemoptysis, pulmonary bleeding; 4. shortness of breath; 5. pain in throat when swallowing; 6. increasing temperature.	 if the correct answer is 1,2 and 3 if the correct answer is 1 and 3 if the correct answer is 2 and 4 if the correct answer is 4 if the correct answer is 1,2,3 and 4 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Choose the correct combination of answers 16. In what cases is it legal to talk about non-resolving pneumonia?	1) by the end of the 4th week from the onset of the disease while the intoxication syndrome persists	high

		2) by the end of the 4th week	
		from the onset of the disease without complete radiological	
		resolution of focal infiltrative	
		changes in the lungs	
		3) by the end of the 2nd week	
		from the onset of the disease	
		without complete radiological	
		resolution of focal infiltrative	
		changes in the lungs4) by the end of the 2nd week	
		from the onset of the disease	
		while the intoxication	
		syndrome persists	
		5) there is no correct answer.	
GC-1 GPC-1	Choose the correct combination of		high
PC-1 PC- 2 PC-7 PC- 11	answers	2) lipase activity	
PC-/ PC- 11	17. Which enzyme level increases in the blood when an ulcer penetrates into the	3) glucose level4) alkaline phosphatase	
	pancreas	activity	
	Parterous	5) none of the above	
GC-1 GPC-1	Draw a conclusion	1) bronchial asthma	high
PC-1 PC- 2	18. A 68-year-old patient with complaints	2) mild COPD	
PC-7 PC- 11	of an unproductive cough in the morning,	3) severe COPD	
	dysphere on exercise, a feeling of shortage	4) everything is correct	
	of air, and fatigue. when studying the function of external respiratory (after	5) pneumonia	
	inhalation of a bronchodilattor) fvc-72%,		
	fev1-35%, fev1/fvc-56%. what is the		
	preliminary diagnosis		
GC-1 GPC-1	Draw a conclusion	A. Correct answers 1, 5 and 9	high
PC-1 PC- 2	19. Patient, 42 years old, undergoing	B. Correct answers are 2, 6	
PC-7 PC- 11	additional check-up. no complaints. from history: he denies any chronic	and 10 B. Correct answers are 3, 7	
	diseases. works as a human resources	and 11	
	manager. work is associated with	D. Correct answers are 4, 8	
	psychoemotional tension. i do not smoke.	and 12	
	stress is relieved in the evening by eating		
	a plenty and prolonged eating while		
	watching tv. does not deny regular alcohol consumption.		
	objective: height 1.76 m, weight 92 kg,		
	from 106 cm. no signs of other somatic		
	pathology detected. probable diagnosis?		
	what is the required scope of		
	investigations to clarify the diagnosis?		
	what treatment should be prescribed? 1. obesity		
	2. metabolic syndrome		
	3. harmful consumption of alcohol		
	4. disordered eating behavior		
	5. calculation of bmi, bioimpedanometry		
	6. laboratory assessment of carbohydrate		
	and lipid metabolism according to criteria		

	 7. rus-audit test 8. consultation with a psychologist and/or clinical psychologist 9. rational low-calorie diet therapy and normalization of physical activity regime 10. metformin 11. refuse to use alcohol 12. cognitive-behavioral psychotherapy 		
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate the correct answer 20. In a 45 year old man, 3 years ago after gastric resection, the content of red cells in the blood is 2.0 x 1012/l; hb 85 g/l; cpu	1) C 2) PP 3) A 4) B6 5) P12	high
	- 1.27. absorption of which vitamin caused this change in erythropoiesis	5) B12	

Test task for diagnostic testing in the discipline:

Discipline name, semester: Outpatient therapy, 11th semester, 6th year

Curriculum	31.05.01
	General Medicine
Qualification	General Medicine
Form of education	Full-time
Department	Internal diseases
Graduating department	Internal diseases

Test task for diagnostic testing in the discipline:

Competence being tested	Exercise	Answer options	Question difficulty type
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Please indicate one correct answer. 1. To diagnose urolithiasis during pregnancy, it is most advisable to use the following research methods 1. overview radiography 2. excretory urography 3. cystoscopy 4. ultrasound 	 1.true 1, 2, 3 2.true 1, 2 3.all of the above are true 4. true 4 all of the above is incorrect 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate one correct answer.</i>2. Which medications rarely lead to constipation	 diuretics aluminum-containing antacids 	low

GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate one correct answer.</i> 3. Define hypostenuria	 calcium preparations antibacterial iron preparations увеличение частоты мочеиспускания reducing the amount of urine reducing the frequency of urination decrease in the specific gravity of urine increase in the specific gravity of urine 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate one correct answer.</i>4. During the day, patient p. excreted800 ml of urine, and at night 1,350.What is the name of this condition?	 dysuria pollakiuria nicturia polyuria oliguria 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Please indicate one correct answer.</i> 5. Which drug should not be used to relieve a hypertensive crisis?	 corinthian enalapril clonidine captopril furosemide 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Please indicate one correct answer. 6 The absolute indication for termination of pregnancy in case of vomiting of pregnant women is: 1. renal insufficiency; 2. liver failure; 3. comatose state; 4. significant reduction in body weight 	 The answers are correct 2, 3. The answers are correct 3. The answers are correct 4. Only the answer is correct 4. All the answers are correct. 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Specify the correct answers</i> 7. Contraindications to oral litholytic therapy for cholelithiasis will be all provisions except	 diarrhea pregnancy exacerbations frequent seizures stone size up to 1.5 cm 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Specify all the correct answers</i> 8. What characterizes the hyperkinetic form of gallbladder dyskinesia	 colicky or cramping pains pain that appears 1-2 hours after eating, usually radiating to the right shoulder blade, collarbone, 	average

		often to the lower back and	
		epigastric region	
		3. With the addition the	
		gallbladder and its	
		accelerated emptying,	
		revealed by X-ray	
		examination	
		4. All of the above is true	
		5. marked weakness	
GC-1 GPC-1	Enter the correct answer	1) there is a predominant	average
PC-1 PC- 2	9. For symptomatic renal arterial	increase in the diastolic	
PC-7 PC-11	hypertension, everything is true except	component of blood	
		pressure	
		-	
		2) kidney angiography will	
		be the optimal diagnostic	
		method	
		3) there is a decrease in	
		plasma renin activity with	
		the administration of ACE	
		4) the captopril sample is	
		positive	
		5) develops more often in	
		old age	
GC-1 GPC-1	Enter the correct answer	1) 3-5 days	average
PC-1 PC- 2	<i>10.</i> What are the approximate terms of	2) 5-15 days	
PC-7 PC-11	temporary disability in angina pectoris 2	3) 10-15 days	
	fc	4) 20 days	
		5) 30-60 days	
GC-1 GPC-1	Specify all the correct answers	1. dark unformed feces with	average
PC-1 PC- 2	11. Specify the characteristics of the	a fetid odor	C
PC-7 PC-11	coprological syndrome in fermentation	2.light foamy feces with a	
	dyspepsia	sour smell	
		3.no starch grains	
		4. yellowish shiny feces	
		5. the presence of neutral	
		fat droplets, fatty acid	
		crystals	
GC-1 GPC-1	Specify all the correct answers	1) IAPF	average
PC-1 PC- 2	12. Which of the drugs may be missing in	2) aspirin	C
PC-7 PC-11	the treatment regimen of a patient with	3) Statin	
	coronary heart disease, angina pectoris 3	4) B-blocker	
	<i>fc and CH 1 ct.</i>	,	
	•	5) cardiac glycosides	
GC-1 GPC-1	Specify all the correct answers	1) reducing the voltage of all	average
PC-1 PC- 2	13. A reliable criterion for myocardial	teeth	
PC-7 PC-11	ischemia in bicycle ergometry is	2) ST segment elevation less	
		than 1 mm	
		3) ST depression of more than	
		2 mm	
		4) the appearance of negative	
		teeth T	
		5) tachycardia	
	-		

GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Specify all the correct answers 14. Which of the following groups of antihypertensive drugs is not allowed prescribe to a patient with hypertension of which bronchial asthma is concomitant	 ACE inhibitors – renitek, kapoten calcium antagonists – corinfar retard, norvask non–selective beta - blockers – obsidan, anaprilin diuretics – hypothyazid, arifon all groups are contraindicated 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Specify all the correct answers 15. Which of the symptoms is most reliable for angina pectoris	 chest pain during exercise rhythm disturbance depression of the st interval, recorded during an attack of pain pathological tooth q ST elevation at the height of pain 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Choose the right combination of answers</i> 16. What condition is characterized by severe, cramping pain in the area of the penis and in the epigastrium, the duration of the attack is 1-2 minutes, with a frequency of 3-10 minutes, vomiting, stools and gas retention, a soft abdomen on palpation, an empty ampoule of the rectum during rectal examination	 acute appendicitis acute cholecystitis intestinal obstruction acute pancreatitis everything is true 	high
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Choose the right combination of answers 17. What is the indication for hemodialysis in a patient with diabetes mellitus complicated by diabetic nephropathy with the development of chronic renal failure?	Iomerular filtration rate lessthan 30 ml/min, serumcreatininelevel>300mmol/l2) glomerular filtration rateless than 15 ml/min, serumcreatininelevel>600mmol/l3) glomerular filtration rateless than 30 ml/min, serumcreatininelevel>300mmol/l4) glomerular filtration rateless than 30 ml/min, serumcreatininelevel>600mmol/l4) glomerular filtration rateless than 30 ml/min, serumcreatininelevel>600mmol/l5) there is no correct answer	high
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Make a conclusion</i> 18. Which symptom is not typical for crf	 1. dry mouth 2. nausea, vomiting 3. itchy skin 4. hemoptysis 5. polyuria 	high

GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Choose the right combination of answers</i> 19. With a combination of hypertension and diabetic nephropathy, which drug group should be included in therapy	 B-blockers IAPF calcium antagonists diuretics a combination of diuretics and ACE inhibitors is necessary 	high
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Enter the correct answer 20. After returning from the restaurant at night, a 53-year-old patient had sharp pains in the area of the thumb of the left foot, the temperature was 38.3 ° c. On examination, swelling of the i metatarsophalangeal joint was revealed, bright hyperemia of the skin around it. Touching the finger causes a sharp increase in pain. What is the most likely diagnosis?	 1) rheumatoid arthritis 2) reactive arthritis 3) gouty arthritis 4) rheumatoid arthritis 5) there is no correct answer 	high

Test task for diagnostic testing in the discipline:

Discipline name, semester: Outpatient therapy, 12th semester, 6th year

Curriculum	31.05.01
	General Medicine
Qualification	General Medicine
Form of education	Full-time
Department	Internal diseases
Graduating department	Internal diseases

Test assignment for diagnostic testing in the discipline:

Проверяема	Задание	Варианты ответов	Тип
я			сложно
компетенци			сти

Я			вопрос а
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate one correct answer. A local general practitioner was called to a 15-year-old teenager due to sudden profuse vomiting and abdominal pain. He got sore throat a week ago. At my grandmother's insistence, i took aspirin for several days. A few hours ago i drank 2 cups of fresh coffee and felt general weakness, dizziness, "flickering of flies". The day before, i ate jelly, fried chicken, chocolate cake, oranges, pineapple, drank tea, pepsi-cola. The vomit resembles coffee grounds, contains lumps of food eaten the day before. The patient is pale, heart rate is 100 in 1 min, weak filling, blood pressure is 100/50 mmhg. The abdomen is soft, painful on palpation in the epigastric region. There was no stool in the last 24 hours. The most likely diagnosis	 a) food poisoning b) acute pancreatitis c) Mallory—Weiss syndrome d) allergic enterocolitis e) erosive gastritis 	a low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Please indicate one correct answer. 2. Preventive treatment of patients with peptic ulcer disease 	 a) it is prescribed to all patients with YAB b) it is prescribed only in cases of risk factors c) preventive treatment is carried out as a seasonal anti-relapse d) a complex of drugs is used during preventive measures e) only H2-histaminoreceptor blockers are prescribed prophylactically 	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate one correct answer. 3. A 49-year-old patient underwent surgery for ulcerative bleeding with perforation 5 years ago. He notes pains similar to the previous ones, ulcerative, but localized in the epigastric region on the left. Notes vomiting at the height of pain and weight loss. Conservative treatment is ineffective. In feces, the gregersen reaction (). Which of the following diseases should be suspected first	a) hypoglycemia syndrome b) agastric asthenia c) adductor loop syndrome d) peptic ulcer of anastomosis e) dumping syndrome	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Please indicate one correct answer. 4. A 52-year-old patient has had a feeling of heaviness in the right hypochondrium, nausea, and slight jaundice for a long time against the background of asthenic complaints. At this time, the biochemical parameters of liver function are impaired: 	 a) portal cirrhosis of the liver b) chronic persistent hepatitis c) biliary cirrhosis of the liver d) chronic active hepatitis e) pseudocyrrhosis of the liver 	low

	alt, bilirubin, sulemic test, thymol test, protein fractions. Relapses are associated		
	with infections. Performance is maintained. which of the following diagnoses corresponds to the clinical syndrome and laboratory data		
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Please indicate one correct answer. 5. A 75-year-old man experiences severe pain in the mesogastric region in combination with vomiting and bloating about 30 minutes after eating. Over the past few months, he has lost 6 kg, as he limited his food intake, fearing that it would provoke the appearance of pain. During the asymptomatic period, an examination of the abdomen without features. noise is heard above the right femoral artery, peripheral pulsation is reduced in both lower extremities. Stool test for latent blood is negative. Fibrogastroduodenoscopy, irrigoscopy and colonoscopy revealed no pathological changes. this condition is most likely due to	a)psychogenic factors b) neoplasm c) inflammation d) ischemia e) transient obstruction	low
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Enter the correct answer 6 The absolute indication for termination of pregnancy in case of vomiting of pregnant women is: 1. renal insufficiency; 2. liver failure; 3. comatose state; 4. significant reduction in body weight 	 The answers are correct 1, 2, 3. The answers are correct 1, 3. The answers are correct 2, 4. Only the answer is correct 4. All the answers are correct. 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11		 diarrhea pregnancy exacerbations frequent seizures Stone size up to 1.5 cm 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Specify all the correct answers 8. What characterizes the hyperkinetic form of gallbladder dyskinesia	 colicky or cramping pains pain that appears 1-2 hours after eating, usually radiating to the right shoulder blade, collarbone, often to the lower back and epigastric region	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Enter the correct answer 9. For symptomatic renal arterial hypertension, everything is true except	1) there is a predominant increase in the diastolic component of blood pressure	average

GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Enter the correct answer</i> 10. What are the approximate terms of temporary disability in angina pectoris 2 fc	 2) kidney angiography will be the optimal diagnostic method 3) there is a decrease in plasma renin activity with the administration of ACE 4) The captopril sample is positive 5) develops more often in old age 1) 3-5 days 2) 5-15 days 3) 10-15 days 4) 20 days 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Enter the correct answer</i> 11. The most informative objective clinical sign for the diagnosis of a generalized form of bronchiectasis is	 5) 30-60 days a) shortening of the percussion tone b) hard breathing c) dry wheezing d) small-bubbly wet wheezes e) localized moist medium- and large-bubbly wheezes 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	<i>Specify all the correct answers</i> 12. An early clinical symptom of central lung cancer is:	 dry cough or cough with mucous sputum, sometimes containing streaks of blood pulmonary hemorrhage shortness of breath and weight loss fever hemoptysis 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Specify all the correct answers 13. The detection of erosions in the antrum of the stomach during fibrogastroduodenoscopy is typical for the treatment of which is carried out 1. granulomatous gastritis 2. menetrieu diseases 3. autoimmune gastritis 4. helicobacter infection 5. gcs 6. proton pump inhibitors 7. gastroprotectors 8. eradication therapy 9. udhk	A. Correct answers 1, 6 and 7 B Correct answers 2, 7 and 9	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	Specify all the correct answers 14. Acute laryngitis in dynamics threatens the development of:	 persistent dysphonia; acute laryngeal stenosis; asthmatic bronchitis; hypersensitivity of the larynx. 	average
GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11	 Specify all the correct answers 15.For the diagnosis of chronic gastritis, pgkazanes: 1. it is enough to collect complaints and anamnesis 2. ultrasound of the obp; 3. morphological examination of the gastric mucosa (soj); 4. identification of hp; 	A. Correct answers 1, and 4 B. Correct answers 3, 4, and 5 C. Correct answers 2, 4 and 6 D. Correct answers 1, 2 and 4	average

	5.esophagogastroduodenoscopy (egds);		
	6. ph-metric of gastric contents.		
GC-1 GPC-1	Choose the right combination of answers	1) by the end of the 4th week from	high
PC-1 PC- 2	16. In which cases is it legitimate to talk	the onset of the disease, while	
PC-7 PC-11	about non-resolving pneumonia	maintaining the intoxication	
		syndrome	
		2) by the end of the 4th week from	
		the onset of the disease without	
		complete X-ray resolution of focal	
		infiltrative changes in the lungs	
		3) by the end of the 2nd week from	
		the onset of the disease without	
		complete X-ray resolution of focal	
		infiltrative changes in the lungs	
		4) by the end of the 2nd week from	
		the onset of the disease, while	
		maintaining the intoxication	
		syndrome	
		5) There is no correct answer.	
GC-1 GPC-1		1. dry mouth	high
PC-1 PC- 2	17. Which symptom is not typical for crf	2. nausea, vomiting	
PC-7 PC- 11		3. itchy skin	
		4. hemoptysis	
		5. polyuria	
GC-1 GPC-1	Choose the right combination of answers	1) In-BLGK.	high
PC-1 PC- 2	18. With a combination of hypertension	2) IAPF	
PC-7 PC-11	and diabetic nephropathy, which drug	3) calcium antagonists	
	group should be included in therapy	4) diuretics	
		5)a combination of diuretics and	
		ACE inhibitors is necessary	
GC-1 GPC-1	Please indicate one correct answer.	a) at least 6 months a year for 2	high
PC-1 PC- 2	19. The WHO criteria for chronic	consecutive years	
PC-7 PC-11	bronchitis is the duration of cough	b) more than 4 months in a given	
		year	
		c) at least 3 months a year for 2 years	
		in a row	
		d) at least 2 months a year for 3 years	
		in a row	

GC-1 GPC- 1 PC-1 PC- 2 PC-7 PC- 11		 rheumatoid arthritis reactive arthritis gouty arthritis rheumatoid arthritis there is no correct answer 	high
--	--	---	------